

Explicit Solutions to the Unitarity Relations for Isovector Nucleon Form Factors

M. I. Krivoruchenko

Institute for Theoretical and Experimental Physics, B.Chermushkinskaya 25

117259 Moscow, Russia

Abstract

The explicit one-dimensional integral representations for isovector nucleon form factors are constructed by solving the unitarity relations in terms of the pion form factor and imaginary parts of the t -channel p -wave πN -scattering amplitudes. The possibility of computation of the high-energy parts of the integrals with the use of the information on the low-energy real parts of the πN -scattering amplitudes is discussed.

1 Introduction

The unitarity relations for isovector nucleon form factors have been discovered by Frazer and Fulco¹ in 1960 yr. These relations were discussed intensively in the 60-th and 70-th years. At present, new precise data are available at low and intermediate momentum transfers both at spacelike and timelike regions (see *e.g.* Ref.2 and references therein). It is desirable therefore to analyze more carefully theoretical methods and current models. In Sect.2, we report new analytical results concerning the unitarity relations. In Sect.3, these results are used to develop an alternative approach for numerical evaluation of the nucleon form factors from the unitarity relations.

2 Solutions to the Unitarity Relations

The unitarity relations for isovector nucleon form factors have the form¹

$$ImF_{iv}(t) = -\frac{k^3}{\sqrt{t}}F_{\pi}^*(t)\Gamma_i(t) \quad (2.1)$$

where $k = \sqrt{t/4 - \mu^2}$, μ is the pion mass, $F_{\pi}(t)$ is the pion form factor, and $\Gamma_i(t)$ are combinations of the t -channel p -wave projections of the πN -scattering amplitudes. The form factors are normalized by $F_{1v}(0) = 1/2$ and $F_{2v}(0) = (\mu_p - \mu_n - 1)/2$ with μ_p and μ_n being the proton and neutron magnetic moments (in n.m.). The proton charge is set equal to unity.

In the approximation of neglecting all but the two-pion intermediate state, the values $\Gamma_i(t)$ obey the once subtracted dispersion relations

$$\Gamma_i(t)/F_{\pi}(t) = Re\frac{\Gamma_i(\kappa)}{F_{\pi}(\kappa)} + \frac{t - \kappa}{\pi}P \int_{-\infty}^a \frac{Im\Gamma_i(t')}{(t' - \kappa)(t' - t)} \frac{dt'}{F_{\pi}(t')}. \quad (2.2)$$

with $a = 4\mu^2 - \mu^4/m^2$. In the FFGS model^{1,3}, the following relation takes place above the two-pion threshold

$$\frac{k^3}{\sqrt{t}}|F_{\pi}(t)|^2 = D(0)\frac{k_{\rho}^3}{m_{\rho}^2\Gamma_{\rho}}\frac{t_0 + t}{t_0}ImF_{\pi}(t + i0). \quad (2.3)$$

where $D(0)$ is the value of the D -function at the origin. The exponentially large parameter t_0 is a unique root of equation $D(-t_0) = 0$, k_{ρ} is the pion momentum at $t = m_{\rho}^2$, m_{ρ} and Γ_{ρ} are the ρ -meson mass and width. In the narrow width limit, $D(0) = m_{\rho}^2$, $t_0 = \infty$.

Combining Eqs.(2.1), (2.2) and (2.3), the once subtracted dispersion integral for the nucleon form factors can be evaluated to give

$$F_{iv}(t) = F_{iv}(-t_0) - D(0)\frac{k_{\rho}^3}{m_{\rho}^2\Gamma_{\rho}}\frac{t_0 + t}{t_0}F_{\pi}(t) \\ (Re\frac{\Gamma_i(\kappa)}{F_{\pi}(\kappa)} + \frac{1}{\pi}P \int_{-\infty}^a \frac{Im\Gamma_i(t')}{(t' - \kappa)} \frac{(t - \kappa)/F_{\pi}(t') - (t' - \kappa)/F_{\pi}(t)}{(t' - t)} dt'). \quad (2.4)$$

For isovector kaon form factor, similar representation has been constructed within the differential approximation scheme⁴. For computation of the nucleon form factors, it is sufficient to know the imaginary parts $Im\Gamma_i(t)$ at the interval $(-\infty, a)$ only.

The imaginary part of the ratio $ImF_{iv}(t)/F(t)$ has a simple form

$$Im\{(F_{iv}(t) - F_{iv}(-t_0))/F_{\pi}(t)\}/\frac{k^3}{\sqrt{t}} = -\frac{1}{\pi} \int_{-\infty}^a \frac{Im\Gamma_i(t')}{t' - t} dt'. \quad (2.5)$$

3 High-Energy Part of the Integrals

The integrals in Eqs.(2.4) and (2.5) are determined by the low- and high-energy parts of the values $Im\Gamma_i(t)$. The values $\Gamma_i(t)$ can be decomposed to the Born and rescattering

parts: $\Gamma_i(t) = \Gamma_{iB}(t) + \Gamma_{iR}(t)$. The values $\Gamma_{iR}(t)$ are known at the interval $-26\mu^2 < t < 0$ from the phase-shift analysis of the πN -scattering amplitudes. The high-energy rescattering parts $\Gamma_{iR}(t)$ at $-\infty < t < -26\mu^2 = b$ are unknown. The problem consists in evaluation of the high-energy parts with the help of the information on the low-energy real parts $Re\Gamma_{iR}(t)$.

In the ratio $ReF_{iv}(t)/F_\pi(t)$, the convergence of the integral can be improved by taking a weighted sum

$$Re\{(F_{iv}(t) - F_{iv}(-t_0))/F_\pi(t)\} = -D(0)\frac{k_\rho^3}{m_\rho^2\Gamma_\rho}\frac{t_0 + t}{t_0}$$

$$\sum_{j=1}^N c_{ij} \left(Re \frac{\Gamma_i(\kappa_j)}{F_\pi(\kappa_j)} + \frac{1}{\pi} P \int_{-\infty}^a \frac{Im\Gamma_i(t')}{(t' - \kappa_j)} \frac{(t - \kappa_j)/F_\pi(t') - (t' - \kappa_j)Re\{1/F_\pi(t)\}}{(t' - t)} dt' \right). \quad (3.1)$$

with the coefficients $\sum_{j=1}^N c_{ij} = 1$. The subtraction points κ_j are assumed to belong to the interval $-26\mu^2 < \kappa_j < 0$. The coefficients c_{ij} must be chosen such as to minimize contributions to the integral from the high-energy part $-\infty < t < -26\mu^2$.

The convergence of the integral in Eq.(2.5) can be improved in the similar way. The right hand side of Eq.(2.5) can be rewritten as follows

$$\sum_{j=1}^N d_{ij} Re \frac{\Gamma_i(\kappa_j)}{F_\pi(\kappa_j)} - \frac{1}{\pi} \left(\int_{-\infty}^b + P \int_b^a \right) Im\Gamma_i(t') \left(\sum_{j=1}^N d_{ij} \frac{1}{t' - \kappa_j} \frac{1}{F_\pi(t')} + \frac{1}{t' - t} \right) dt' \quad (3.2)$$

The weight coefficients d_{ij} must be chosen such as to suppress the high-energy part of the integral. To improve convergence of the integral, one should require $\sum_{j=1}^N d_{ij} = 0$.

4 Conclusion

In the standard approach¹, one should compute two integrals numerically. The first one comes from the dispersion integral for the $\Gamma_i(t)$. The second one comes from the dispersion integral for the nucleon form factors. We showed that the last integral can be evaluated analytically. In doing so, we arrived to the explicit one-dimensional integral representations (2.4) for the nucleon form factors in terms of the pion form factor and the amplitudes $\Gamma_i(t)$ at $t < a$. These representations provide an alternative possibility for numerical analysis of the problem.

References

- [1] W.R.Frazer and J.Fulco, Phys.Rev.Lett. **2** (1959) 365; Phys.Rev. **115** (1960) 1763; **117** (1960) 1609.

- [2] T.A.Armstrong et al., Phys.Rev.Lett. **70** (1993) 1212.
- [3] G.Gounaris and J.J.Sakurai, Phys.Rev.Lett. **21** (1968) 244.
- [4] M.I.Krivoruchenko, Pis'ma ZhETF **58** (1993) 7; Z.Phys. **A350** (1995) 343.